

Orbifolds are not commutative geometries

Adam Rennie^{†*} and Joseph C. Várilly^{‡||*}

[†]Institute for Mathematical Sciences, University of Copenhagen
Universitetsparken 5, DK-2100 Copenhagen, Denmark

[‡]Departamento de Física Teórica I, Universidad Complutense,
Madrid 28040, Spain

^{||}Departamento de Matemáticas, Universidad de Costa Rica,
2060 San José, Costa Rica

21 March 2007

Abstract

In this note we show that the crucial orientation condition for commutative geometries fails for the natural spectral triple of an orbifold M/G .

1 Introduction

Recently, in [4], it was shown that a spectral triple $(\mathcal{A}, \mathcal{H}, \mathcal{D})$ with $\mathcal{A}$ commutative and unital is the Dirac triple of a spin^c manifold $(C^\infty(M), L^2(M, S), \not{D})$ if and only if $(\mathcal{A}, \mathcal{H}, \mathcal{D})$ satisfies certain additional conditions. These conditions are a mild strengthening and generalisation of those proposed in [2]. (A spectral triple with $\mathcal{A}$ commutative and satisfying some version of these conditions could thus be called a commutative geometry, hence the title).

In this note we show precisely why orbifolds of the form M/G , with G a finite group acting by isometries on M , fail to satisfy these conditions. For such orbifolds we can define a commutative spectral triple, and though we do not show it here, these spectral triples do satisfy the majority of those conditions. However, the crucial orientation condition, which provides a volume form, tangent bundle and ultimately gives the local coordinates, fails for the spectral triples of the orbifolds.

After a brief recollection of the conditions, we go directly to the demonstration of the failure of the orientation condition. To finish, we briefly comment on more general orbifolds.

2 Geometric conditions for commutative spectral triples

In [4] we have shown that under a small number of postulates, mildly extending those of [2], any spectral triple $(\mathcal{A}, \mathcal{H}, \mathcal{D})$ whose algebra $\mathcal{A}$ is *commutative* and unital satisfies $\mathcal{A} = C^\infty(M)$ where M is a smooth closed manifold carrying a spin structure, $\mathcal{H}$ is its L^2 -spinor space and $\mathcal{D}$ is a Dirac operator plus an endomorphism of the spinor bundle.

*email: rennie@math.ku.dk, varilly@cariari.ucr.ac.cr

Recall that a (unital) spectral triple consists of a unital algebra $\mathcal{A}$ faithfully represented on a Hilbert space $\mathcal{H}$, and a selfadjoint operator $\mathcal{D}$ on $\mathcal{H}$, with compact resolvent, such that $\mathcal{A}$ preserves its domain $\text{Dom } \mathcal{D}$ and $[\mathcal{D}, a]$ extends to a bounded operator on $\mathcal{H}$ for each $a \in \mathcal{A}$.

Assume furthermore that $\mathcal{A}$ is commutative. Denote by A its norm completion in $\mathcal{B}(\mathcal{H})$ and by $M = \text{sp}(A) = \text{sp}(\mathcal{A})$ its character space (Gelfand spectrum), which is assumed to be a separable C^* -algebra. The following postulates, of an algebraic or operatorial nature, are needed for the reconstruction of the manifold, spin structure and Riemannian metric.

1. *Dimension*: For some $p \in \{1, 2, 3, \dots\}$, the operator $\langle \mathcal{D} \rangle^{-1} := (1 + \mathcal{D}^2)^{-1/2}$ lies in $\mathcal{L}^{p, \infty}$ and $\langle \mathcal{D} \rangle^{-p} \in \mathcal{L}^{1, \infty}$ has positive Dixmier traces: $\text{Tr}_\omega \langle \mathcal{D} \rangle^{-p} > 0$ for all ω .

When p is even, $\mathcal{H}$ is $\mathbb{Z}_2$ -graded by a selfadjoint unitary Γ commuting with $\mathcal{A}$ and anticommuting with $\mathcal{D}$. When p is odd, $\mathcal{H}$ is ungraded and we put $\Gamma = 1$ for convenience.

2. *Regularity*: $(\mathcal{A}, \mathcal{H}, \mathcal{D})$ is a QC^∞ in the sense of [1]: if $\delta(x) := [|\mathcal{D}|, x]$ for $x \in \mathcal{B}(\mathcal{H})$, then $\mathcal{A} \cup [\mathcal{D}, \mathcal{A}] \subseteq \bigcap_{m=1}^\infty \text{Dom } \delta^m$. Moreover, $\mathcal{A}$ is complete in the locally convex topology given by the seminorms $q_m(a) := \|\delta^m a\|$ and $q'_m(a) := \|\delta^m([\mathcal{D}, a])\|$, for $m = 0, 1, 2, \dots$.
3. *Finiteness*: The prehilbert space $\mathcal{H}_\infty = \bigcap_{m \geq 1} \text{Dom } \mathcal{D}^m$ is a finitely generated projective $\mathcal{A}$ -module.
4. *Absolute continuity*: If $a > 0$ in $\mathcal{A}$, then $\text{Tr}_\omega a \langle \mathcal{D} \rangle^{-p} > 0$ for all ω .
5. *First order*: $[[\mathcal{D}, a], b] = 0$ for $a, b \in \mathcal{A}$.
6. *Orientability*: There is a Hochschild p -cycle

$$\mathbf{c} = \sum_{\alpha=1}^n a_\alpha^0 \otimes a_\alpha^1 \otimes \dots \otimes a_\alpha^p \in Z_p(\mathcal{A}, \mathcal{A}) \quad (2.1a)$$

which is represented on $\mathcal{H}$ by Γ , that is,

$$\pi_{\mathcal{D}}(\mathbf{c}) \equiv \sum_{\alpha=1}^n a_\alpha^0 [\mathcal{D}, a_\alpha^1] \dots [\mathcal{D}, a_\alpha^p] = \Gamma. \quad (2.1b)$$

7. *Closedness*: $\text{Tr}_\omega (\Gamma [\mathcal{D}, a_1] \dots [\mathcal{D}, a_p] \langle \mathcal{D} \rangle^{-p}) = 0$ for all $a_1, \dots, a_p \in \mathcal{A}$.
8. *Spin^c or Morita equivalence*: The C^* -module completion of $\mathcal{H}_\infty$ is a Morita equivalence bimodule between A and the norm completion of $\mathcal{C}_{\mathcal{D}}(\mathcal{A})$.
9. *Spin or Reality*: There is an antiunitary operator J on $\mathcal{H}$ such that $Ja^*J^{-1} = a$ for all $a \in \mathcal{A}$; $J^2 = \pm 1$; $J\mathcal{D}J^{-1} = \pm \mathcal{D}$; and $J\Gamma J^{-1} = \pm \Gamma$ if p is even; with the same signs (depending only on $p \bmod 8$) as occur when $\mathcal{A} = C^\infty(M)$ with M a p -dimensional spin manifold, $\mathcal{D}$ is a Dirac operator and J is the charge conjugation.
10. *Connectivity*: There is an orthogonal family of projectors $p_j \in \mathcal{A}$ such that $\text{Id}_{\mathcal{A}} = \sum_j p_j$ and

$$(a \in \mathcal{A} \text{ with } [\mathcal{D}, a] = 0) \iff a = \sum_j \lambda_j p_j \text{ for some } \{\lambda_j\} \subset \mathbb{C}.$$

For a detailed discussion of how each of these conditions contributes to the reconstruction of the smooth manifold M , we refer to [4]. For instance, the orientability condition plays a role in ensuring that all operators appearing under Dixmier traces above are “measurable”, which means that it does not matter which Dixmier trace is used. Another point to notice is that, by the Serre–Swan theorem, there is a complex vector bundle $S \rightarrow M$ (that turns out to be the spinor bundle) for which $\mathcal{H}_\infty \subset \Gamma(M, S)$, the latter being the A -module of continuous sections of S .

Naturally, the conclusion that M is a smooth manifold is a rather strong one, and one may wonder whether some “almost-manifolds” could also be allowed. For instance, one can build a spectral triple on an orbifold, satisfying many, but not all, of the above postulates. While one normally uses Lie groupoids [3] to describe orbifolds, which entails noncommutative algebras, one could construct a spectral triple on a quotient space M/G using the algebra $\mathcal{A} = C^\infty(M)^G$ of smooth invariant functions, which satisfies many of the properties listed above. We next show that this attempt fails, precisely on account of the orientability condition.

3 A G -invariant spectral triple

Consider the following example of a commutative spectral triple: let (M, g) be a compact boundaryless smooth Riemannian manifold with metric g of dimension p , with a spin structure (so that in particular M is orientable), and let G be a finite group acting by isometries on M . We shall suppose that M is connected; if not, it has finitely many components and we may restrict our attention to the subgroup of G that preserves a given component. Assume that this action lifts to an action of G on the spinor bundle $S \rightarrow M$, for which the Dirac operator $\mathcal{D}_g$ is G -invariant. Now let $\mathcal{A} := C^\infty(M)^G$ be the algebra of G -invariant smooth functions on M , $\mathcal{H} := L^2(M, S)^G$ be the Hilbert space of G -invariant L^2 -spinors, and let $\mathcal{D}$ be the restriction of $\mathcal{D}_g$ to a selfadjoint operator on $\mathcal{H}$. Then $(\mathcal{A}, \mathcal{H}, \mathcal{D})$ is a spectral triple over a commutative unital algebra.

(It might happen that an isometric action of G on M lifts only to that of an extension $\tilde{G}$ of G by $\mathbb{Z}_2$, acting by automorphisms of the spinor bundle. Since $\tilde{G}$ is also a finite group, the arguments below are not materially affected by this variant; so we may as well assume that the isometric action of G lifts directly to the spinor bundle.)

We claim that this example *cannot satisfy the orientation condition* unless G acts freely on M .

If the action of G on M is free, then the orbit space $X = M/G$ is a smooth Riemannian manifold and this spectral triple may be identified with a Dirac spectral triple over $C^\infty(X)$. In particular, all the defining conditions listed in [4] will hold automatically for $(\mathcal{A}, \mathcal{H}, \mathcal{D})$. However, if the action is not free, then M/G is instead an orbifold. We do not need to consider this quotient space as such, but much of the standard terminology of orbifold theory is useful. We follow the notation of [3], for the most part.

The isotropy subgroup of $x \in M$ is $G_x := \{h \in G : h \cdot x = x\} \leq G$, and the fixed set of $h \in G$ is $\Sigma_h := \{x \in M : h \cdot x = x\} \subset M$. The *singular locus* is

$$\Sigma_G := \bigcup_{h \neq 1} \Sigma_h = \{x \in M : G_x \neq 1\},$$

so that G acts freely on its complement. The exponential map at $x \in M$ coming from the metric g gives a diffeomorphism $\exp_x : U \rightarrow W$ from a small open neighbourhood of $0 \in T_x M$

to a neighbourhood W of x in M . If $h \in G$, then dh_x is an orthogonal linear transformation from $T_x M$ to $T_{h \cdot x} M$, such that

$$\exp_{h \cdot x} \circ dh_x = h \circ \exp_x : U \rightarrow h \cdot W, \quad (3.1a)$$

When $h \in G_x$, we may take W to be G_x -invariant and may assume that each dh_x preserves U , so that

$$\exp_x \circ dh_x = h \circ \exp_x : U \rightarrow W \quad (3.1b)$$

in that case. Note [3, Lemma 2.10] that if $h \in G_x$, then $\{y \in \Sigma_h : dh_y = 1 \text{ on } T_y M\}$ is both closed and open in M by (3.1a), so that if M is connected, then $h \mapsto dh_x : G_x \rightarrow \text{Aut}(T_x M)$ is injective for all $x \in M$. Together with (3.1b), this shows that Σ_h has empty interior if $h \neq 1$. Thus the singular locus Σ_G is a closed subset of M with empty interior.

By transposition, each dh_x acts on the cotangent spaces, taking $T_{h \cdot x}^* M$ to $T_x^* M$, again orthogonally with respect to the transposed metric g^{-1} . In other words, dh has a right action on covectors.

Any G -invariant function f equals its average over any cyclic subgroup of G :

$$f(x) = \frac{1}{|h|} \sum_{j=0}^{|h|-1} f(h^j \cdot x),$$

where $|h|$ denotes the order of h . If $h \in G_x$, taking commutators with the Dirac operator gives

$$[\mathcal{D}, f](x) = c(df(x)) = \frac{1}{|h|} \sum_{j=0}^{|h|-1} c(df(x) \lhd dh_x^j),$$

where c denotes Clifford multiplication. Note that (3.1b) implies that the notation dh_x^j is unambiguous: $\exp_x \circ (dh_x)^j = h^j \circ \exp_x = \exp_x \circ (dh^j)_x$, so that $(dh_x)^j = (dh^j)_x$ on $T_x M$.

Let $x \in \Sigma_G$. The Cartan–Dieudonné theorem, showing that every orthogonal transformation on $T_x M$ is the product of at most p reflections, allows us to choose, for each $h \in G_x$, an orthonormal basis for $T_x M$ such that

$$dh_x = \text{diag}(R_{\theta_1}, R_{\theta_2}, \dots, R_{\theta_k}, \pm 1, \dots, \pm 1),$$

where each R_{θ_j} is a two-by-two rotation matrix (itself a product of two reflections).

Consider first the case where G acts on M by orientation-preserving isometries, so that each dh_x is a rotation in $SO(T_x M)$. Then we can write

$$[\mathcal{D}, f](x) = c(df(x)) = c(u) + c(v_1) + \dots + c(v_k),$$

where $u \in T_x^* M$ is invariant under dh_x and $v_r \lhd dh_x = R_{-\theta_r} v_r$, for $r = 1, \dots, k$. Then

$$\begin{aligned} [\mathcal{D}, f](x) &= \frac{1}{|h|} \sum_{j=0}^{|h|-1} c(df(x) \lhd dh_x^j) \\ &= \frac{1}{|h|} \sum_{j=0}^{|h|-1} c(u + R_{-\theta_1}^j v_1 + \dots + R_{-\theta_k}^j v_k) \\ &= c(u) + \frac{1}{|h|} \sum_{r=1}^k \left(\sum_{j=0}^{|h|-1} R_{-\theta_r}^j \right) v_r = c(u), \end{aligned}$$

since $\sum_{j=0}^{|h|-1} R_{-\theta_r}^j = 0$: that follows since the order of each rotation $R_{-\theta_r}$ divides $|h|$.

Hence, if f is any G -invariant function, $[\mathcal{D}, f](x)$ is fixed by the rotation dh_x and so it is (Clifford multiplication by) a covector perpendicular to all the planes of rotation of h .

Therefore, if $G_x \neq 1$ and if $a_0, a_1, \dots, a_p$ are G -invariant functions, then $a_0 [\mathcal{D}, a_1] \dots [\mathcal{D}, a_n]$ has vanishing skewsymmetrisation at x :

$$\frac{1}{p!} \sum_{\sigma \in S_p} (-1)^\sigma \sum_{\alpha} a_0(x) [\mathcal{D}, a_{\sigma(1)}](x) \dots [\mathcal{D}, a_{\sigma(p)}](x) = 0, \quad (3.2)$$

since, under the usual linear isomorphism $\mathbb{C}\ell(T_x^* M, g_x) \simeq \Lambda^\bullet T_x^* M$, the left hand side corresponds to a p -covector in $\Lambda^p V$, where $V = \bigcap_{h \in G_x} \ker(\lhd dh_x)$ has dimension less than p .

In particular, on inserting the entries of the Hochschild p -cycle (2.1a) in Eqn. (3.2), we obtain

$$\Gamma'(x) = \frac{1}{p!} \sum_{\sigma \in S_p} (-1)^\sigma \sum_{\alpha} a_{\alpha}^0(x) [\mathcal{D}, a_{\alpha}^{\sigma(1)}](x) \dots [\mathcal{D}, a_{\alpha}^{\sigma(p)}](x) = 0 \quad (3.3)$$

for all $x \in \Sigma_G$. We have written Γ' for the skewsymmetrized version of the represented cycle (2.1b). Here Γ' may be regarded as an endomorphism of the $\mathcal{A}$ -module $\mathcal{H}_{\infty}$ and thereby as a continuous section in $\Gamma(M, \text{End } S)$.

Next consider the case where some G_x contains a reflection.

Let $x \in \Sigma_G$ and $h \in G_x$ be such that dh_x is a reflection in the hyperplane $v^\perp$, for some unit covector $v \in T_x^* M$. Thus, if f is G -invariant, then

$$\begin{aligned} c(df(x)) &= [\mathcal{D}, f](x) = \frac{1}{2}(c(df(x)) + c(df(x) \lhd dh_x)) \\ &= \frac{1}{2}c(df(x)) + \frac{1}{2}(c(df(x)) - 2g_x(v, df(x))c(v)) \\ &= c(df(x)) - g_x(v, df(x))c(v). \end{aligned}$$

Thus $g_x(v, df(x)) = 0$, so $df(x)$ is a covector in the hyperplane $v^\perp$. Again, any p such covectors have null exterior product, so that $\Gamma'(x) = 0$ in this case also.

Suppose now that there are finitely many G -invariant functions $\{a_{\alpha}^j : j = 0, \dots, p, \alpha = 1, \dots, n\}$ such that $\mathbf{c}$ as given by (2.1a) is a Hochschild p -cycle, and that Equation (2.1b) holds with Γ being proportional to Clifford multiplication by the volume element, as happens when $\mathcal{D}$ is the Dirac operator. By using local orthonormal bases of 1-forms, one can check that the skewsymmetrization of this section of the Clifford algebra bundle is, up to a constant, Clifford multiplication by the Riemannian volume form vol_g on M . As such, the skewsymmetrization is nowhere vanishing and the norm of $\Gamma'(y) \in \text{End } S_y$ has a positive lower bound. Since we have shown that for G -invariant functions there is always a neighbourhood of the singular locus Σ_G on which $\|\Gamma'(y)\|$ is smaller than any such lower bound, the orientability condition cannot be satisfied.

The argument extends to any compact orbifold, since these are locally of the form M/G with G finite. Indeed, such an orbifold X is a topological space that may be covered by finitely many orbifold charts [3], each of the form U/G where U is an open connected subset of $\mathbb{R}^p$ and G is a finite group of diffeomorphisms of U ; one may impose a Riemannian metric on U for which G acts by isometries. Overlapping charts may use different finite groups, but all are linked by compatibility homomorphisms. The singular locus of the orbifold is the union of finitely many sets Σ_G in each such chart, and the above argument shows that the candidate

for a volume form, if it comes from a G -invariant p -form on each U , must always vanish on the singular locus, and therefore cannot satisfy (2.1b).

While the argument is local, it is not clear that for a general orbifold we have a good candidate for a (commutative) algebra of functions: this is highly dependent on the pasting conditions [3]. When dealing with quotients M/Γ , where Γ is an infinite discrete group acting locally freely on M (that is, all isotropy subgroups are finite), there is such a candidate algebra but no simple method of suitably encoding the local freedom. It is therefore more appropriate to employ the groupoid algebra in these situations.

Acknowledgements

We thank Gianni Landi for first asking us about the admissibility of orbifolds as commutative geometries. Later, others have raised the same question. This work was supported by an ARC grant, DP0211367, by the Statens Naturvidenskabelige Forskningsråd, Denmark. Support from the University Complutense de Madrid and the Vicerrectoría de Investigación of the Universidad de Costa Rica is acknowledged. JCV is grateful to Ryszard Nest for kind hospitality at the University of Copenhagen.

References

- [1] A. L. Carey, J. Phillips, A. Rennie and F. A. Sukochev, “The Hochschild class of the Chern character for semifinite spectral triples”, *J. Funct. Anal.* **213** (2004), 111–153.
- [2] A. Connes, “Gravity coupled with matter and foundation of noncommutative geometry”, *Commun. Math. Phys.* **182** (1996), 155–176.
- [3] I. Moerdijk and J. Mrčun, *Introduction to Foliations and Lie Groupoids*, Cambridge University Press, Cambridge, 2003.
- [4] A. Rennie and J. C. Várilly, “Reconstruction of manifolds in noncommutative geometry”, math.oa/0610418 v3, Copenhagen, 2006.